

Delphion

RESEARCH

INTEGRATED IAM

SERVICES

INSIDE DELPHION

[Log Out](#) [Work Files](#) [Saved Searches](#)[My Account](#) | [Products](#) | [News](#) | [Events](#)Search: [Quick/Number](#) [Boolean](#) [Advanced](#)[Help](#)

## The Delphion Integrated View


Buy N w: [More choices...](#)Tools: Add to Work File: [Create new Work File](#) Vi w: [INPADOC](#) | Jump to: [Top](#) ☐ [Email this to a friend](#)Title: **JP2000164249A2: NONAQUEOUS ELECTROLYTE AND NONAQUEOUS ELECTROLYTE SECONDARY BATTERY**Country: **JP Japan**Kind: **A2 Document Laid open to Public inspection**Inventor: **HAYASHI TAKASHI;  
TAN HIROAKI;**Assignee: **MITSUI CHEMICALS INC**  
[News, Profiles, Stocks and More about this company](#)Published / Filed: **June 16, 2000 / Nov. 27, 1998**Application Number: **JP1998000336632**IPC Code: **H01M 10/40;**Priority Number: **Nov. 27, 1998 JP1998000336632**

Abstract:

PROBLEM TO BE SOLVED: To provide an electrolyte with low heat generation speed and high safety by constituting with a nonaqueous solvent containing fluorine-containing cyanoethyl ether compound and an electrolyte.

SOLUTION: A nonaqueous electrolyte contains fluorine-containing cyanoethyl ether compound represented by the formula: X-(OR)<sub>n</sub>-OCH<sub>2</sub>CH<sub>2</sub>-CN. In the formula, X shows 1-10C fluorine atom substituted hydrocarbon group, R shows 2-4C alkylene group, and (n) is 0-30. The nonaqueous solvent preferably contains cyclic carbonate and/or chain carbonate, the cyclic carbonate is a compound containing 2-5C alkylene group, and the chain carbonate is a compound containing 1-5C hydrocarbon group. An electrolyte is selected from LiPF<sub>6</sub>, LiBF<sub>6</sub>, and LiOSO<sub>2</sub>R<sub>1</sub>. R<sub>1</sub> shows 1-6C perfluoroalkyl group. A negative electrode containing metallic lithium, a lithium-containing alloy, and a carbon material capable of doping/undoping lithium ions as a negative active material is used.

COPYRIGHT: (C)2000,JPO

Family: **None**Other Abstract Info: **CHEMABS 133(02)020143S CHEMABS 133(02)020143S DERABS G2000-456796 DERABS G2000-456796**[View Image](#)

1 page

[Nominate](#)[this for the Gallery...](#)



(19)

(11) Publication number: **2000164249 A**

Generated Document.

**PATENT ABSTRACTS OF JAPAN**(21) Application number: **10336632**(51) Intl. Cl.: **H01M 10/40**(22) Application date: **27.11.98**

(30) Priority:

(43) Date of application  
publication: **16.06.00**(84) Designated contracting  
states:(71) Applicant: **MITSUI CHEMICALS INC**(72) Inventor: **HAYASHI TAKASHI  
TAN HIROAKI**

(74) Representative:

**(54) NONAQUEOUS  
ELECTROLYTE AND  
NONAQUEOUS  
ELECTROLYTE  
SECONDARY BATTERY**

(57) Abstract:

**PROBLEM TO BE SOLVED:** To provide an electrolyte with low heat generation speed and high safety by constituting with a nonaqueous solvent containing fluorine-containing cyanoethyl ether compound and an electrolyte.

**SOLUTION:** A nonaqueous electrolyte contains fluorine-containing cyanoethyl ether compound represented by the formula:  $X-(OR)_n-OCH_2CH_2-CN$ . In the formula, X shows 1-10C fluorine atom substituted hydrocarbon group, R shows 2-4C alkylene group, and (n) is 0-30. The nonaqueous solvent preferably contains cyclic carbonate and/or chain carbonate, the cyclic carbonate is a compound containing 2-5C alkylene group, and the chain carbonate is a compound containing 1-5C hydrocarbon group. An electrolyte is selected from  $LiPF_6$ ,  $LiBF_6$ , and  $LiOSO_2R_1$ .  $R_1$  shows 1-6C

perfluoroalkyl group. A negative  
~~electrode containing metallic lithium, a~~  
lithium-containing alloy, and a carbon  
material capable of doping/undoping  
lithium ions as a negative active  
material is used.

COPYRIGHT: (C)2000,JPO